

- mia: a multivariate analysis from the randomized GCLLSG CLL8 trial. *J Clin Oncol.* 2012;30(9):980-988.
20. Fischer K, Bahlo J, Fink AM, et al. Long-term remissions after FCR chemoimmunotherapy in previously untreated patients with CLL: updated results of the CLL8 trial. *Blood.* 2016;127(2):208-215.
 21. Fischer K, Cramer P, Busch R, et al. Bendamustine in combination with rituximab for previously untreated patients with chronic lymphocytic leukemia: a multicenter phase II trial of the German Chronic Lymphocytic Leukemia Study Group. *J Clin Oncol.* 2012;30(26):3209-3216.
 22. Eichhorst B, Fink AM, Bahlo J, et al. First-line chemoimmunotherapy with bendamustine and rituximab versus fludarabine, cyclophosphamide, and rituximab in patients with advanced chronic lymphocytic leukaemia (CLL10): an international, open-label, randomised, phase 3, non-inferiority trial. *Lancet Oncol.* 2016;17(7):928-942.
 23. Kutsch N, Bahlo J, Robrecht S, et al. Long Term Follow-up Data and Health-Related Quality of Life in Frontline Therapy of Fit Patients Treated With FCR Versus BR (CLL10 Trial of the GCLLSG). *Hemasphere.* 2020;4(1):e336.
 24. Goede V, Fischer K, Busch R, et al. Obinutuzumab plus chlorambucil in patients with CLL and coexisting conditions. *N Engl J Med.* 2014;370(12):1101-1110.
 25. Goede V, Fischer K, Engelke A, et al. Obinutuzumab as frontline treatment of chronic lymphocytic leukemia: updated results of the CLL11 study. *Leukemia.* 2015;29(7):1602-1604.
 26. Mulligan SP, Gill DS, Turner P, et al. A Randomised Dose De-Escalation Safety Study of Oral Fludarabine, (+/-) Oral Cyclophosphamide and Intravenous Rituximab (OFOCIR) As First-Line Therapy of Fit Patients with Chronic Lymphocytic Leukemia (CLL) Aged >=65 Years - End of Recruitment Analysis of Response and Toxicity of the Australasian Leukaemia and Lymphoma Group (ALLG) and CLL Australian Research Consortium (CLLARC) CLL5 Study. *ASH Annual Meeting Abstracts.* 2012;120(21):436.
 27. Smolej L, Brychtova Y, Cmun E, et al. Low-dose fludarabine and cyclophosphamide combined with rituximab in the first-line treatment of elderly/comorbid patients with chronic lymphocytic leukaemia/small lymphocytic lymphoma (CLL/SLL): long-term results of project Q-lite by the Czech CLL Study Group. *Br J Haematol.* 2021;193(4):769-778.
 28. Burger JA. Inhibiting B-cell receptor signaling pathways in chronic lymphocytic leukemia. *Curr Hematol Malig Rep.* 2012;7(1):26-33.
 29. Byrd JC, Brown JR, O'Brien S, et al. Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med.* 2014;371(3):213-223.
 30. Munir T, Brown JR, O'Brien S, et al. Final analysis from RESONATE: Up to six years of follow-up on ibrutinib in patients with previously treated chronic lymphocytic leukemia or small lymphocytic lymphoma. *Am J Hematol.* 2019;94(12):1353-1363.
 31. Burger JA, Tedeschi A, Barr PM, et al. Ibrutinib as Initial Therapy for Patients with Chronic Lymphocytic Leukemia. *N Engl J Med.* 2015;373(25):2425-2437.
 32. Woyach JA, Ruppert AS, Heerema NA, et al. Ibrutinib Regimens versus Chemoimmunotherapy in Older Patients with Untreated CLL. *N Engl J Med.* 2018;379(26):2517-2528.
 33. Shanafelt TD, Wang XV, Kay NE, et al. Ibrutinib-Rituximab or Chemoimmunotherapy for Chronic Lymphocytic Leukemia. *N Engl J Med.* 2019;381(5):432-443.
 34. Shanafelt TD, Wang XV, Hanson CA, et al. Long-term outcomes for ibrutinib-rituximab and chemoimmunotherapy in CLL: updated results of the E1912 trial. *Blood.* 2022;140(2):112-120.
 35. Mato AR, Nabhan C, Thompson MC, et al. Toxicities and outcomes of 621 ibrutinib-treated chronic lymphocytic leukemia patients in the United States: a real-world analysis. *Haematologica.* 2018;103(5):874-879. doi: 10.3324/haematol.2017.182907. Epub 2018 Feb 1. PMID: 29419429; PMCID: PMC5927982.
 36. Paydas S. Management of adverse effects/toxicity of ibrutinib. *Crit Rev Oncol Hematol.* 2019;136:56-63.
 37. Lasica M, Tam CS. Management of Ibrutinib Toxicities: a Practical Guide. *Curr Hematol Malig Rep.* 2020;15(3):177-186.
 38. Stephens DM, Byrd JC. How I manage ibrutinib intolerance and complications in patients with chronic lymphocytic leukemia. *Blood.* 2019;133(12):1298-1307.
 39. Burger JA, Barr PM, Robak T, et al. Long-term efficacy and safety of first-line ibrutinib treatment for patients with CLL/SLL: 5 years of follow-up from the phase 3 RESONATE-2 study. *Leukemia.* 2020;34(3):787-798.
 40. Seymour JF, Byrd JC, Ghia P, et al. Detailed safety profile of acalabrutinib vs ibrutinib in previously treated chronic lymphocytic leukemia in the ELEVATE-RR trial. *Blood.* 2023;142(8):687-699.
 41. Burger JA, Sivina M, Jain N, et al. Randomized trial of ibrutinib vs ibrutinib plus rituximab in patients with chronic lymphocytic leukemia. *Blood.* 2019;133(10):1011-1019.
 42. Kater AP, Owen C, Moreno C, et al. Fixed-Duration Ibrutinib-Venetoclax in Patients with Chronic Lymphocytic Leukemia and Comorbidities. *NEJM Evid.* 2022;1(7).
 43. Danilov AV, Persky DO. Incorporating acalabrutinib, a selective next-generation Bruton tyrosine kinase inhibitor, into clinical practice for the treatment of haematological malignancies. *Br J Haematol.* 2021;193(1):15-25.
 44. Thompson PA, Burger JA. Bruton's tyrosine kinase inhibitors: first and second generation agents for patients with Chronic Lymphocytic Leukemia (CLL). *Expert Opin Investig Drugs.* 2018;27(1):31-42.
 45. Ghia P, Pluta A, Wach M, et al. ASCEND: Phase III, Randomized Trial of Acalabrutinib Versus Idelalisib Plus Rituximab or Bendamustine Plus Rituximab in Relapsed or Refractory Chronic Lymphocytic Leukemia. *J Clin Oncol.* 2020;38(25):2849-2861.
 46. Sharman JP, Egyed M, Jurczak W, et al. Acalabrutinib with or without obinutuzumab versus chlorambucil and obinutuzumab for treatment-naïve chronic lymphocytic leukaemia (ELEVATE TN): a randomised, controlled, phase 3 trial. *Lancet.* 2020;395(10232):1278-1291.
 47. Byrd JC, Hillmen P, Ghia P, et al. Acalabrutinib Versus Ibrutinib in Previously Treated Chronic Lymphocytic Leukemia: Results of the First Randomized Phase III Trial. *J Clin Oncol.* 2021;39(31):3441-3452.
 48. Brown JR, Eichhorst B, Hillmen P, et al. Zanubrutinib or Ibrutinib in Relapsed or Refractory Chronic Lymphocytic Leukemia. *N Engl J Med.* 2023;388(4):319-332.
 49. Furman RR, Sharman JP, Coutre SE, et al. Idelalisib and rituximab in relapsed chronic lymphocytic leukemia. *N Engl J Med.* 2014;370(11):997-1007.
 50. Flinn IW, Hillmen P, Montillo M, et al. The phase 3 DUO trial: duvelisib vs ofatumumab in relapsed and refractory CLL/SLL. *Blood.* 2018;132(23):2446-2455.
 51. Sharman JP, Coutre SE, Furman RR, et al. Final Results of a Randomized, Phase III Study of Rituximab With or Without Idelalisib Followed by Open-Label Idelalisib in Patients With Relapsed Chronic Lymphocytic Leukemia. *J Clin Oncol.* 2019;37(16):1391-1402.
 52. Spacek M, Smolej L, Simkovic M, et al. Idelalisib plus rituximab versus ibrutinib in the treatment of relapsed/refractory chronic lymphocytic leukaemia: A real-world analysis from the Chronic Lymphocytic Leukemia Patients Registry (CLLE-AR). *Br J Haematol.* 2023;202(1):40-47.
 53. Ryan CE, Davids MS. BCL-2 Inhibitors, Present and Future. *Cancer J.* 2019;25(6):401-409.
 54. Stilgenbauer S, Eichhorst B, Schetelig J, et al. Venetoclax in relapsed or refractory chronic lymphocytic leukaemia with 17p deletion: a multicentre, open-label, phase 2 study. *Lancet Oncol.* 2016;17(6):768-778.
 55. Roberts AW, Davids MS, Pagel JM, et al. Targeting BCL2 with Venetoclax in Relapsed Chronic Lymphocytic Leukemia. *N Engl J Med.* 2016;374(4):311-322.
 56. Cairo MS, Bishop M. Tumour lysis syndrome: new therapeutic strategies and classification. *Br J Haematol.* 2004;127(1):3-11.
 57. Howard SC, Jones DP, Pui CH. The tumor lysis syndrome. *N Engl J Med.* 2011;364(19):1844-1854.
 58. Gribben JG. Practical management of tumour lysis syndrome in venetoclax-treated patients with chronic lymphocytic leukaemia. *Br J Haematol.* 2020;188(6):844-851.
 59. Seymour JF, Kipps TJ, Eichhorst B, et al. Venetoclax-Rituximab in Relapsed or Refractory Chronic Lymphocytic Leukemia. *N Engl J Med.* 2018;378(12):1107-1120.
 60. Fischer K, Al-Sawaf O, Bahlo J, et al. Venetoclax and Obinutuzumab in Patients with CLL and Coexisting Conditions. *N Engl J Med.* 2019;380(23):2225-2236.
 61. Rawstron AC, Fazi C, Agathangelidis A, et al. A complementary role of multiparameter flow cytometry and high-throughput sequencing for minimal residual disease detection in chronic lymphocytic leukemia: an European Research Initiative on CLL study. *Leukemia.* 2016;30(4):929-936.
 62. Kater AP, Seymour JF, Hillmen P, et al. Fixed Duration of Venetoclax-Rituximab in Relapsed/Refractory Chronic Lymphocytic Leukemia Eradicates Minimal Residual Disease and Prolongs Survival: Post-Treatment Follow-Up of the MURANO Phase III Study. *J Clin Oncol.* 2019;37(4):269-277.
 63. Al-Sawaf O, Zhang C, Tandon M, et al. Venetoclax plus obinutuzumab versus chlorambucil plus obinutuzumab for previously untreated chronic lymphocytic leukaemia (CLL14): follow-up results from a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* 2020;21(9):1188-1200.
 64. Dreger P. Is There a Role for Cellular Therapy in Chronic Lymphocytic Leukemia? *Cancer J.* 2021;27(4):297-305.
 65. Tournilhac O, van Gelder M, Eikema DJ, et al. The European landscape on allogeneic haematopoietic cell transplantation in Chronic Lymphocytic Leukaemia between 2009 and 2019: a perspective from the Chronic Malignancies Working Party of the EBMT. *Bone Marrow Transplant.* 2023;58(6):621-624.
 66. Mancikova V, Smida M. Current State of CART-T-Cell Therapy in Chronic Lymphocytic Leukemia. *Int J Mol Sci.* 2021;22(11).
 67. Thompson MC, Harrup RA, Coombs CC, et al. Venetoclax retreatment of patients with chronic lymphocytic leukemia after a previous venetoclax-based regimen. *Blood Adv.* 2022;6(15):4553-4557.
 68. Ahn IE, Underbayev C, Albitar A, et al. Clonal evolution leading to ibrutinib resistance in chronic lymphocytic leukemia. *Blood.* 2017;129(11):1469-1479.
 69. Seymour JF, Kipps TJ, Eichhorst BF, et al. Enduring undetectable MRD and updated outcomes in relapsed/refractory CLL after fixed-duration venetoclax-rituximab. *Blood.* 2022;140(8):839-850.
 70. Shanafelt TD, Borah BJ, Finnes HD, et al. Impact of ibrutinib and idelalisib on the pharmaceutical cost of treating chronic lymphocytic leukemia at the individual and societal levels. *J Oncol Pract.* 2015;11(3):252-258.
 71. Hilal T, Betcher JA, Leis JF. Economic Impact of Oral Therapies for Chronic Lymphocytic Leukemia-the Burden of Novelty. *Curr Hematol Malig Rep.* 2018;13(4):237-243.
 72. Davids MS, Chatterjee A, Ravelo A, et al. Cost-Effectiveness of a 12-Month Fixed Duration of Venetoclax in Combination with Obinutuzumab in First-Line Chronic Lymphocytic Leukemia in the United States. *Blood.* 2019;134(Supplement_1):4741.
 73. Patel K, Isufi I, Kothari S, et al. Cost-Effectiveness of First-Line Ibrutinib versus Third-Line in Patients with Untreated Chronic Lymphocytic Leukemia. *Blood.* 2020;136(17):1946-1955.
 74. Simkovic M, Vodarek P, Motyckova M, et al. Rituximab, Cyclophosphamide and Dexamethasone (RCD) Chemoimmunotherapy for Relapsed Chronic Lymphocytic Leukaemia. *Eur J Clin Invest.* 2021;51(4):e13421.
 75. Smolej L, Špaček M, Pospíšilová, et al. Doporučení pro diagnostiku a léčbu chronické lymfocytární leukemie 2021. *Transfuze Hematol. dnes* 2021;27(1):91-106.
 76. Eichhorst B, Niemann CU, Kater AP, et al. First-Line Venetoclax Combinations in Chronic Lymphocytic Leukemia. *N Engl J Med.* 2023;388(19):1739-1754.
 77. Mato AR, Woyach JA, Brown JR, et al. Pirtobrutinib after a Covalent BTK Inhibitor in Chronic Lymphocytic Leukemia. *N Engl J Med.* 2023;389(1):33-44.