

LITERATURA

1. Ferlay J, Ervik M, Colombet M, et al. Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>. Published 2022. Accessed May 19, 2024.
2. GBD 2017 Stomach Cancer Collaborators. The global, regional, and national burden of stomach cancer in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease study 2017. *Lancet Gastroenterol Hepatol*. 2020;5(1):42–54. doi: 10.1016/S2468-1253(19)30328-0. Epub 2019 Oct 21. Erratum in: *Lancet Gastroenterol Hepatol*. 2020;5(3):e2. PMID: 31648970; PMCID: PMC7033564.
3. Colquhoun A, Arnold M, Ferlay J, et al. Global patterns of cardia and non-cardia gastric cancer incidence in 2012. *Gut*. 2015;64(12):1881–1888. doi: 10.1136/gutjnl-2014-308915. Epub 2015 Mar 6. PMID: 25748648.
4. Pimentel-Nunes P, Dinis-Ribeiro M, Ponchon T, et al. Endoscopic submucosal dissection: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy*. 2015;47(9):829–854. doi: 10.1055/s-0034-1392882. Epub 2015 Aug 28. PMID: 26317585.
5. Lordick F, Carneiro F, Cascinu S, et al. Gastric cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol*. 2022;33(10):1005–1020. doi: 10.1016/j.annonc.2022.07.004. Epub 2022 Jul 29. PMID: 35914639.
6. Cunningham D, Allum WH, Stenning SP, et al. Perioperative chemotherapy versus surgery alone for resectable gastroesophageal cancer. *N Engl J Med*. 2006;355(1):11–20. doi: 10.1056/NEJMoa055531. PMID: 16822992.
7. Ychou M, Boige V, Pignon JP, et al. Perioperative chemotherapy compared with surgery alone for resectable gastroesophageal adenocarcinoma: an FNCLCC and FFCD multicenter phase III trial. *J Clin Oncol*. 2011;29(13):1715–1721. doi: 10.1200/JCO.2010.33.0597. Epub 2011 Mar 28. PMID: 21444866.
8. Alderson D, Cunningham D, Nankivell M, et al. Neoadjuvant cisplatin and fluorouracil versus epirubicin, cisplatin, and capecitabine followed by resection in patients with oesophageal adenocarcinoma (UK MRC OE05): an open-label, randomised phase 3 trial. *Lancet Oncol*. 2017;18(9):1249–1260. doi: 10.1016/S1470-2045(17)30447-3. Epub 2017 Aug 4. PMID: 28784312; PMCID: PMC5585417.
9. Al-Batran SE, Homann N, Pauligk C, et al. Perioperative chemotherapy with fluorouracil plus leucovorin, oxaliplatin, and docetaxel versus fluorouracil or capecitabine plus cisplatin and epirubicin for locally advanced, resectable gastric or gastro-oesophageal junction adenocarcinoma (FLOT4): a randomised, phase 2/3 trial. *Lancet*. 2019;393(10184):1948–1957. doi: 10.1016/S0140-6736(18)32557-1. Epub 2019 Apr 11. PMID: 30982686.
10. Smyth EC, Fassan M, Cunningham D, et al. Effect of Pathologic Tumor Response and Nodal Status on Survival in the Medical Research Council Adjuvant Gastric Infusional Chemotherapy Trial. *J Clin Oncol*. 2016;34(23):2721–2727. doi: 10.1200/JCO.2015.65.7692. Epub 2016 Jun 13. PMID: 27298411; PMCID: PMC5019747.
11. Rahman S, Thomas B, Maynard N, et al. Impact of postoperative chemotherapy on survival for oesophagogastric adenocarcinoma after preoperative chemotherapy and surgery. *Br J Surg*. 2022;109(2):227–236. doi: 10.1093/bjs/znab427. PMID: 34910129; PMCID: PMC10364695.
12. Smyth E, Mauer M, Cella C, et al. O-6 EORTC 1707 VESTIGE: Adjuvant immunotherapy in patients (pts) with resected gastroesophageal adenocarcinoma (GEA) following preoperative chemotherapy with high risk for recurrence (ypN+ and/or R1) an open-label randomized controlled phase II study. *Annals of Oncology*. 2023;34:S182–S183. <https://doi.org/10.1016/j.annonc.2023.04.021>.
13. Cats A, Jansen EPM, van Grieken NCT, et al. Chemotherapy versus chemoradiotherapy after surgery and preoperative chemotherapy for resectable gastric cancer (CRITICS): an international, open-label, randomised phase 3 trial. *Lancet Oncol*. 2018;19(5):616–628. doi: 10.1016/S1470-2045(18)30132-3. Epub 2018 Apr 9. PMID: 29650363.
14. de Steur WO, van Amelsfoort RM, Hartgrink HH, et al. Adjuvant chemotherapy is superior to chemoradiation after D2 surgery for gastric cancer in the per-protocol analysis of the randomized CRITICS trial. *Ann Oncol*. 2021;32(3):360–367. doi: 10.1016/j.annonc.2020.11.004. Epub 2020 Nov 20. PMID: 33227408.
15. Shapiro J, van Lanschot JJB, Hulshof MCCM, et al. Neoadjuvant chemoradiotherapy plus surgery versus surgery alone for oesophageal or junctional cancer (CROSS): long-term results of a randomised controlled trial. *Lancet Oncol*. 2015;16(9):1090–1098. doi: 10.1016/S1470-2045(15)00040-6. Epub 2015 Aug 5. PMID: 26254683.
16. Eyck BM, van Lanschot JJB, Hulshof MCCM, et al. Ten-Year Outcome of Neoadjuvant Chemoradiotherapy Plus Surgery for Esophageal Cancer: The Randomized Controlled CROSS Trial. *J Clin Oncol*. 2021;39(18):1995–2004. doi: 10.1200/JCO.20.03614. Epub 2021 Apr 23. PMID: 33891478.
17. Reynolds JV, Preston SR, O'Neill B, et al. Trimodality therapy versus perioperative chemotherapy in the management of locally advanced adenocarcinoma of the oesophagus and oesophagogastric junction (Neo-AEGIS): an open-label, randomised, phase 3 trial. *Lancet Gastroenterol Hepatol*. 2023;8(11):1015–1027. doi: 10.1016/S2468-1253(23)00243-1. Epub 2023 Sep 18. PMID: 37734399; PMCID: PMC10567579.
18. Hoepfner J, Brunner T, Lordick F, et al. Prospective randomized multicenter phase III trial comparing perioperative chemotherapy (FLOT protocol) to neoadjuvant chemoradiation (CROSS protocol) in patients with adenocarcinoma of the esophagus (ESOPEC trial). *JCO*. 2024;42:LBA1-LBA1. doi: 10.1200/JCO.2024.42.17_suppl.LBA1.
19. Bang YJ, Kim YW, Yang HK, et al. Adjuvant capecitabine and oxaliplatin for gastric cancer after D2 gastrectomy (CLASSIC): a phase 3 open-label, randomised controlled trial. *Lancet*. 2012;379(9813):315–321. doi: 10.1016/S0140-6736(11)61873-4. Epub 2012 Jan 7. PMID: 22226517.
20. Noh SH, Park SR, Yang HK, et al. Adjuvant capecitabine plus oxaliplatin for gastric cancer after D2 gastrectomy (CLASSIC): 5-year follow-up of an open-label, randomised phase 3 trial. *Lancet Oncol*. 2014;15(12):1389–1396. doi: 10.1016/S1470-2045(14)70473-5. Epub 2014 Oct 15. PMID: 25439693.
21. Sakuramoto S, Sasako M, Yamaguchi T, et al. Adjuvant chemotherapy for gastric cancer with S-1, an oral fluoropyrimidine. *N Engl J Med*. 2007;357(18):1810–1820. doi: 10.1056/NEJMoa072252. Erratum in: *N Engl J Med*. 2008;358(18):1977. PMID: 17978289.
22. Sasako M, Sakuramoto S, Katai H, et al. Five-year outcomes of a randomized phase III trial comparing adjuvant chemotherapy with S-1 versus surgery alone in stage II or III gastric cancer. *J Clin Oncol*. 2011;29(33):4387–4393. doi: 10.1200/JCO.2011.36.5908. Epub 2011 Oct 17. PMID: 22010012.
23. Kakeji Y, Yoshida K, Kodera Y, et al. Three-year outcomes of a randomized phase III trial comparing adjuvant chemotherapy with S-1 plus docetaxel versus S-1 alone in stage III gastric cancer: JACCRO GC-07. *Gastric Cancer*. 2022;25(1):188–196. doi: 10.1007/s12020-021-01224-2. Epub 2021 Aug 5. PMID: 34351555.
24. GASTRIC (Global Advanced/Adjuvant Stomach Tumor Research International Collaboration) Group; Paoletti X, Oba K, Burzykowski T, Michiels S, Ohashi Y, Pignon JP, Rougier P, Sakamoto J, Sargent D, Sasako M, Van Cutsem E, Buyse M. Benefit of adjuvant chemotherapy for resectable gastric cancer: a meta-analysis. *JAMA*. 2010;303(17):1729–1737. doi: 10.1001/jama.2010.534. PMID: 20442389.
25. Terasima M, Kang YK, Kim YW, et al. ATTRACTION-5: A phase 3 study of nivolumab plus chemotherapy as postoperative adjuvant treatment for pathological stage III (pStage III) gastric or gastroesophageal junction (G/GJE) cancer. *JCO*. 2023;41:4000–4000. doi: 10.1200/JCO.2023.41.16_suppl.4000.
26. Lee J, Lim DH, Kim S, et al. Phase III trial comparing capecitabine plus cisplatin versus capecitabine plus cisplatin with concurrent capecitabine radiotherapy in completely resected gastric cancer with D2 lymph node dissection: the ARTIST trial. *J Clin Oncol*. 2012;30(3):268–273. doi: 10.1200/JCO.2011.39.1953. Epub 2011 Dec 19. PMID: 22184384.
27. Park SH, Lim DH, Sohn TS, et al. A randomized phase III trial comparing adjuvant single-agent S1, S-1 with oxaliplatin, and postoperative chemoradiation with S-1 and oxaliplatin in patients with node-positive gastric cancer after D2 resection: the ARTIST 2 trial*. *Ann Oncol*. 2021;32(3):368–374. doi: 10.1016/j.annonc.2020.11.017. Epub 2020 Dec 3. PMID: 33278599.
28. Macdonald JS, Smalley SR, Benedetti J, et al. Chemoradiotherapy after surgery compared with surgery alone for adenocarcinoma of the stomach or gastroesophageal junction. *N Engl J Med*. 2001;345(10):725–730. doi: 10.1056/NEJMoa010187. PMID: 11547741.
29. Smalley SR, Benedetti JK, Haller DG, et al. Updated analysis of SWOG-directed intergroup study 0116: a phase III trial of adjuvant radiochemotherapy versus observation after curative gastric cancer resection. *J Clin Oncol*. 2012;30(19):2327–2333. doi: 10.1200/JCO.2011.36.7136. Epub 2012 May 14. PMID: 22585691; PMCID: PMC4517071.
30. Dikken JL, Jansen EP, Cats A, et al. Impact of the extent of surgery and postoperative chemoradiotherapy on recurrence patterns in gastric cancer. *J Clin Oncol*. 2010;28(14):2430–2436. doi: 10.1200/JCO.2009.26.9654. Epub 2010 Apr 5. PMID: 20368551.
31. Bang YJ, Van Cutsem E, Feyereislova A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet*. 2010;376(9742):687–697. doi: 10.1016/S0140-6736(10)61121-X. Epub 2010 Aug 19. Erratum in: *Lancet*. 2010;376(9749):1302. PMID: 20728210.
32. Hofheinz RD, Hegewisch-Becker S, Kunzmann V, et al. Trastuzumab in combination with 5-fluorouracil, leucovorin, oxaliplatin and docetaxel as perioperative treatment for patients with human epidermal growth factor receptor 2-positive locally advanced esophagogastric adenocarcinoma: A phase II trial of the Arbeitsgemeinschaft Internistische Onkologie Gastric Cancer Study Group. *Int J Cancer*. 2021;149(6):1322–1331. doi: 10.1002/ijc.33696. Epub 2021 May 29. PMID: 34019698.
33. Hofheinz RD, Merx K, Haag GM, et al. FLOT Versus FLOT/Trastuzumab/Pertuzumab Perioperative Therapy of Human Epidermal Growth Factor Receptor 2-Positive Resectable Esophagogastric Adenocarcinoma: A Randomized Phase II Trial of the AIO EGA Study Group. *J Clin Oncol*. 2022;40(32):3750–3761. doi: 10.1200/JCO.22.00380. Epub 2022 Jun 16. PMID: 35709415.
34. Wagner AD, Grabsch HJ, Mauer M, et al. Integration of trastuzumab (T), with or without pertuzumab (P), into perioperative chemotherapy (CT) of HER-2 positive gastric (GC) and esophagogastric junction cancer (EGJC): First results of the EORTC 1203 INNOVATION study, in collaboration with the Korean Cancer Study Group, and the Dutch Upper GI Cancer group. *JCO*. 2023;41:4057–4057. doi: 10.1200/JCO.2023.41.16_suppl.4057.
35. Fuchs CS, Tomasek J, Yong CJ, et al. Ramucicrumab monotherapy for previously treated advanced gastric or gastro-oesophageal junction adenocarcinoma (REGARD): an international, randomised, multicentre, placebo-controlled, phase 3 trial. *Lancet*. 2014;383(9911):31–39. doi: 10.1016/S0140-6736(13)61719-5. Epub 2013 Oct 3. PMID: 24094768.
36. Wilke H, Muro K, Van Cutsem E, et al. Ramucicrumab plus paclitaxel versus placebo plus paclitaxel in patients with previously treated advanced gastric or gastro-oesophageal junction adenocarcinoma (RAINBOW): a double-blind, randomised phase 3 trial. *Lancet Oncol*. 2014;15(11):1224–1235. doi: 10.1016/S1470-2045(14)70420-6. Epub 2014 Sep 17. PMID: 25240821.

**Další literatura u autora
a na www.onkologiecs.cz**